



Planting density affected biomass and grain yield of maize for seed production in an arid region of Northwest China

JIANG Xuelian^{1,2}, TONG Ling^{1*}, KANG Shaozhong¹, LI Fusheng³, LI Donghao¹, QIN Yonghui¹, SHI Rongchao¹, LI Jianbing⁴

¹ Center for Agricultural Water Research in China, China Agricultural University, Beijing 100083, China;

² Key Laboratory of Biochemistry and Molecular Biology in Universities of Shandong, Weifang University, Weifang 261061, China;

³ College of Agriculture, Guangxi University, Nanning 530005, China;

⁴ Environmental Science and Engineering Program, University of Northern British Columbia, Prince George BC V2N 4Z9, Canada

Abstract: Field experiments were conducted from 2012 to 2015 in an arid region of Northwest China to investigate the effects of planting density on plant growth, yield, and water use efficiency (WUE) of maize for seed production. Five planting densities of 6.75, 8.25, 9.75, 11.25 and 12.75 plants/m² were conducted in 2012, and a planting density of 14.25 plants/m² was added from 2013 to 2015. Through comparison with the AquaCrop yield model, a modified model was developed to estimate the biomass accumulation and yield under different planting densities using adjustment coefficient for normalized biomass water productivity and harvest index. It was found that the modified yield model had a better performance and could generate results with higher determination coefficient and lower error. The results indicated that higher planting density increased the leaf area index and biomass accumulation, but decreased the biomass accumulation per plant. The total yield increased rapidly as planting density increased to 11.25 plants/m², but only a slight increase was observed when the density was greater than 11.25 plants/m². The WUE also reached the maximum when planting density was 11.25 plants/m², which was the recommended planting density of maize for seed production in Northwest China.

Keywords: planting density; yield model; biomass accumulation; grain yield; water use efficiency; Northwest China

Citation: JIANG Xuelian, TONG Ling, KANG Shaozhong, LI Fusheng, LI Donghao, QIN Yonghui, SHI Rongchao, LI Jianbing. 2018. Planting density affected biomass and grain yield of maize for seed production in an arid region of Northwest China. *Journal of Arid Land*, 10(2): 292–303. <https://doi.org/10.1007/s40333-018-0098-7>

1 Introduction

Planting density is one of the most important agronomic factors affecting crop grain yield and water use efficiency (WUE). An increased planting density usually improves the seasonal interception of solar radiation (Harper et al., 1979; Papadopoulos and Pararajasingham, 1997; Westgate et al., 1997) that leads to increased canopy photosynthesis and biomass accumulation (Loomis and Connor, 1992; Coetto et al., 2013), and thus a higher grain yield and water productivity would be reached (Lang et al., 1956; Holt and Timmons, 1968; Fulton, 1970; Qiu et al., 2013). However, the

*Corresponding author: TONG Ling (E-mail: tongling2001@cau.edu.cn)

Received 2017-05-22; revised 2017-12-06; accepted 2017-12-22

© Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Science Press and Springer-Verlag GmbH Germany, part of Springer Nature 2018

grain yield tends to decline when planting density is above a certain level (Mohamed, 1999; Griesh and Yakout, 2001) because of limited supply of carbon and nitrogen under intense interplant competition for intercepted radiation, soil nutrient and water. This would also increase the barrenness and decrease the kernel number (Lemcoff and Loomis, 1986). Consequently, an optimal planting density exists for maximizing the utilization of available resources and achieving the maximum yield per unit area by coordinating crop population and individual development. For example, a few studies have reported that maize grain yield and WUE increased with planting density when it was below an optimum level (Lang et al., 1956; Holt and Timmons, 1968; Westgate et al., 1997). The optimal planting density of maize varies depending on maize varieties and microclimate conditions. Olson and Sanders (1988) noted that the optimal maize planting density ranged from 3.00 to over 9.00 plants/m². Under the surface irrigation conditions, the optimal planting density was found from 5.00 to 5.60 plants/m² in Egypt (Mohamed, 1999; Griesh and Yakout, 2001), while El-Hendawy et al. (2008) observed an optimal planting density of 7.10 plants/m² under full irrigation.

In recent years, the planting area of maize has increased rapidly in China, while the demand for maize seeds has also significantly increased. Particularly, in the arid region of Northwest China, the natural isolation conditions, abundant resources of light and heat are quite suitable for maize for seed production. As a result, the maize for seed production has become the main irrigated crop in this region with a rapidly increased planting area. As different from hybrid maize, the maize for seed production includes the female and male parents of short inbred lines without heterosis. A variety of studies have been conducted to examine the effects of planting density on the yield of common maize (Fulton, 1970; Mohamed, 1999; Griesh and Yakout, 2001; El-Hendawy et al., 2008). However, few studies were reported to investigate the effects of planting density on the yield and WUE of maize for seed production, especially by using yield models. The AquaCrop model developed recently by the Food and Agriculture Organization of the United Nations (FAO) assumed that the growth rate of aboveground biomass was linearly proportional to transpiration through biomass water productivity (WP), while the yield was determined by the harvest index (HI) and multiplication of the aboveground biomass (Raes et al., 2009; Steduto et al., 2009). It was found that HI increased as the increase of planting densities but started to decline after the optimal planting density achieved (DeLougherty and Crookston, 1979; Rahmati, 2009). To obtain a conservative estimation for a given crop, we should normalize WP for climate by taking into account the ratio of transpiration to reference crop evapotranspiration (Steduto et al., 2009). For example, the normalized biomass water productivity (WP*) was found to be about 30 to 35 g/m² for C₄ crops (Raes et al., 2011). However, it is unclear whether WP* and HI in the AquaCrop yield model can be taken as constants under different planting densities of maize for seed production, and this needs further investigation. The objectives of this study were then (1) to investigate the effects of planting density on plant growth, yield and WUE of maize for seed production; (2) to modify WP* and HI in the AquaCrop yield model and improve the estimated accuracy under different planting densities; and (3) to determine the recommended planting density of maize for seed production in the arid region of Northwest China.

2 Materials and methods

2.1 Experimental site

The experiments were conducted during the growing seasons from 2012 to 2015 at Shiyanghe Experimental Station of China Agricultural University (37°52'N, 102°50'E; 1581 m a.s.l.) located in Wuwei, Gansu Province, China. This station is in a typical continental temperate climate zone, with the mean annual precipitation of 164.4 mm, mean annual pan evaporation of 2000 mm, average groundwater table lower than 25 m below the ground surface, mean annual sunshine duration over 3028 h, mean frost-free days of 150 d, and annual mean temperature of 8.8°C (Jiang et al., 2014). The soil texture at the experimental station was a light sandy loam with the mean soil dry bulk density of 1.38 g/cm³ and field water capacity of 0.29 cm³/cm³.

2.2 Experimental design

Maize for seed production is normally planted at a planting density of 9.75–10.5 plants/m² in Northwest China. In 2012, the experiments were designed five planting densities, including 6.75, 8.25, 9.75, 11.25 and 12.75 plants/m² referred as D1, D2, D3, D4 and D5, respectively. A planting density of 14.25 plants/m² was added from 2013 to 2015, referred as D6. A randomized complete block design with three replicates per planting density treatment was used. The experimental plot size was in 9.6 m long and 6.0 m wide. Maize in all plots was sown in each one row-male parents with seven-row female parents in 2012, and with five-row female parents from 2013 to 2015. Each planting density treatment had the same row spacing of 40 cm, and thus the six different planting densities of D1, D2, D3, D4, D5 and D6 were associated with different plant spacing within row of 37, 30, 25, 22, 20 and 18 cm, respectively. Before sowing, all plots were fertilized with N of 136 kg/hm², P₂O₅ of 240 kg/hm² and K₂O of 50 kg/hm² as basal fertilizers. After fertilization, the soil surface in each plot was partly covered with 1.2 m width of film and 0.4 m width of bare soil between two rows. The plots were top-dressed with N of 364 kg/hm² on 10 June 2012, 5 June 2013, 8 June 2014, and 8 June 2015, respectively. The irrigation quota was 100 mm using a border irrigation method. Table 1 lists the meteorological variables, irrigation and planting time of maize.

Table 1 Meteorological variable as well as the planting and irrigation time over the whole growth stage of maize for seed production in 2012–2015

Year	Meteorological variable				Planting and irrigation time	
	R _s (W/m ²)	P (mm)	VPD (kPa)	T _a (°C)	Planting	Irrigation
2012	265.38	129.40	1.23	19.02	16 Apr (Female)	6 Jun
					23 Apr (Male)	26 Jun
					26 Apr (Male)	13 Jul
					–	8 Aug
					–	27 Aug
2013	208.73	68.40	1.45	19.29	13 Apr (Female)	6 Jun
					20 Apr (Male)	26 Jun
					23 Apr (Male)	13 Jul
					–	8 Aug
					–	27 Aug
2014	222.06	206.2	1.28	17.94	15 Apr (Female)	1 Jun
					22 Apr (Male)	1 Jul
					25 Apr (Male)	20 Jul
					–	23 Aug
					–	–
2015	228.78	142.00	1.46	19.22	17 Apr (Female)	8 Jun
					24 Apr (Male)	2 Jul
					29 Apr (Male)	21 Jul
					–	7 Aug
					–	26 Aug

Note: R_s, solar radiation; P, precipitation; VPD, vapor pressure deficit; T_a, air temperature; –, no data.

2.3 Measurements

2.3.1 Meteorological variables

The meteorological variables including solar radiation (R_s), precipitation (P), air temperature (T_a) and relative humidity (RH) during the growing seasons of 2012–2015, were continuously observed by using a standard automatic weather station (Hobo, Onset Computer Corp., USA) that was 100 m away from the experimental plot. The 30 min averages of all meteorological variables were calculated and recorded using a data logger.

2.3.2 Leaf area index (LAI) and biomass accumulation

Five female plants in each plot were randomly selected to measure the LAI and biomass

accumulation. Leaf length and maximum width were measured with a ruler at intervals of 7–10 days after 15 days since sowing. The leaf area was determined by summing the rectangular area (length×maximum width) of each completely developed leaf, and then adjusted by a factor of 0.74 (Li et al., 2008). The leaf area per plant was divided by the surface area per plant to obtain LAI. The plant samples were collected at each growing stage. The total biomass accumulation was obtained after the leaves were oven dried at 60°C when the weights of plant samples kept constant in three hours, and then weighed using an electronic scale with the precision of 0.01 g.

2.3.3 Determination of transpiration

Soil water content in the root-zone was monitored by Diviner 2000 system (Sentek Pty Ltd., Australia). Two PVC access tubes were installed below the mulching and bare soil respectively in each plot. The measurements were made at 0.1 m intervals within 0–1 m soil depth every 5–7 days as well as before and after each irrigation and heavy rain event. The calibration of measurements was conducted using gravimetric method as near as possible to the probe.

The evapotranspiration (ET (mm)) was estimated through soil water balance analysis (Rana and Katerji, 2000). Since the experimental plot was flat and the rainfall was not intensive, the surface runoff was neglected. The groundwater recharge was also negligible since groundwater level was lower than 25 m below the ground surface. Moreover, the measured soil water content at the 90–100 cm soil layer did not change before and after each irrigation event, thus the drainage can be ignored. ET is then estimated by Equation 1.

$$ET = P_e + I - \Delta W, \quad (1)$$

where P_e is the effective precipitation (mm), I is the irrigation amount (mm), and ΔW represents the water content change (mm) in the root-zone that is determined by Equation 2.

$$\Delta W = W_{t_2} - W_{t_1}, \quad (2)$$

where W_{t_1} and W_{t_2} represent the mean water content (mm) in the root-zone at time t_1 and t_2 , respectively.

The soil evaporation (E (mm)) was measured with two micro-lysimeters placed within the bare soil between two plastic films in each plot. The micro-lysimeter with 20 cm height was made from PVC tubes with the inner and outer cylinder diameters of 10 and 11 cm, respectively. The outer cylinder was fixed into the soil with its top leveling with soil surface. The inner cylinder inside the outer cylinder was filled with an intact soil core, and was weighed at 19:00 LST every day by an electronic scale with the precision of 0.1 g. After the measurement of E , the crop transpiration under different planting densities during the growing season (T_r (mm)) was calculated by Equation 3.

$$T_r = ET - E. \quad (3)$$

2.3.4 Grain yield, water use efficiency (WUE) and harvest index (HI)

Fifteen plants in the center of each plot were randomly selected for manual harvesting in each season. The seeds were weighed after sun-drying to obtain grain yield (Y , t/hm²). The WUE (kg/m³) was calculated using Equation 4.

$$WUE = \frac{Y}{ET}. \quad (4)$$

The HI (%) under different planting densities was given by Equation 5.

$$HI = \frac{Y}{B_A}, \quad (5)$$

where B_A is the aboveground biomass accumulation (t/hm²).

2.4 Estimation of biomass accumulation and yield

In AquaCrop model, the B_A was estimated using the crop transpiration under different planting densities during the growing season (T_r) under different planting densities during the growing season and normalized water productivity under standard crop management practice (WP_0^*) by Equation 6 (Steduto et al., 2009):

$$B_A = K_{S_b} WP_0^* \sum \frac{T_r}{ET_0}, \quad (6)$$

where ET_0 is the reference crop evapotranspiration calculated according to the FAO Penman-Monteith equation (Allen et al., 1998); K_{S_b} is the air temperature stress coefficient which can be taken as 1 due to no temperature stress occurred during the growing season in this study; WP_0^* is the normalized water productivity under standard crop management practice which was represented by a reference planting density of 11.25 plants/m² in the study area.

The harvestable yield (Y_A (t/hm²)) was then calculated by B_A and HI under standard crop management practice (HI_0) using Equation 7.

$$Y_A = HI_0 B_A. \quad (7)$$

The AquaCrop yield model was modified in this study by introducing the adjusted WP^* under different planting density (WP_{adj}^*) and the adjusted HI under different planting density (HI_{adj}). A linear relationship (Eq. 8) was assumed between the relative WP^* (WP_{adj}^*/WP_0^*) and the relative planting density (D_{adj}/D_0):

$$\frac{WP_{adj}^*}{WP_0^*} = a \frac{D_{adj}}{D_0} + b, \quad (8)$$

where D_{adj} is the planting density (plants/m²) and D_0 is the reference planting density (11.25 plants/m²). a and b represent the empirical coefficient which can be fitted using the measured transpiration and aboveground biomass accumulation in 2012 and 2013, and they were found as 0.37 and 0.61, respectively. HI_{adj} can be calculated as Equation 9.

$$\frac{HI_{adj} - HI_0}{D_{adj} - D_0} = cD_{adj} + d, \quad (9)$$

where c and d represent the empirical coefficients which were fitted using the measured yield and aboveground biomass accumulation in 2012 and 2013, and they were found as -0.18 and 1.29, respectively.

The WP_{adj}^* and HI_{adj} can be calculated by Equations 10 and 11 based on Equations 8 and 9.

$$WP_{adj}^* = \left(a \frac{D_{adj}}{D_0} + b \right) WP_0^*, \quad (10)$$

$$HI_{adj} = HI_0 + (cD_{adj} + d)(D_{adj} - D_0). \quad (11)$$

The aboveground biomass accumulation (B_d) and grain yield (Y_d) under different planting densities were then obtained using the modified model (Eqs. 12 and 13).

$$B_d = \left(a \frac{D_{adj}}{D_0} + b \right) WP_0^* \sum \frac{T_r}{ET_0}, \quad (12)$$

$$Y_d = (HI_0 + (cD_{adj} + d)(D_{adj} - D_0)) B_d. \quad (13)$$

2.5 Data analysis and evaluation of model performance

SPSS 13.0 software (SPSS Inc., USA) was used for statistical analysis. The means were compared using Duncan's multiple-range test at the 5% probability level. The modeling performance was evaluated based on a linear regression between the estimated (E_i) and observed (Q_i) values of yield. Meanwhile, the mean absolute bias error (MAE) and root mean square error (RMSE) were included and were calculated as Equations 14 and 15 (Legates and McCabe, 1999), respectively.

$$MAE = \frac{1}{N} \sum_{i=1}^N |Q_i - E_i|, \quad (14)$$

$$\text{RMSE} = \left\{ \frac{1}{N} \sum_{i=1}^N (E_i - Q_i)^2 \right\}^{1/2}, \quad (15)$$

where N is the number of data samples.

3 Results and discussion

3.1 Effect of planting density on LAI and biomass accumulation

Figure 1 shows the LAI of maize for seed production under different planting densities in all seasons. Under each planting density, the LAI started to increase rapidly at the shooting stage, reached the maximum at the heading stage, and declined at the maturity stage due to leaf yellowing or wilting. During each season, the LAI increased with planting density, except at the seeding stage. The LAI difference between different planting densities reached the maximum at the heading stage. The maximum LAI differences of about 60% in 2012, 70% in 2013, 88% in 2014, and 80% in 2015 were observed between D1 planting density and the maximum planting density, respectively.




Fig. 1 Leaf area index (LAI) of maize for seed production under different planting densities. Error bars denote standard error of the mean. D1, D2, D3, D4, D5 and D6 are planting densities of 6.75, 8.25, 9.75, 11.25, 12.75 and 14.25 plants/m², respectively.

Figures 2 and 3 present the variation of biomass accumulation and the biomass per plant with planting densities in different years. It was observed that the biomass accumulation and biomass per plant showed a logistic growth pattern, and increased gradually with the advance of plant growth regardless of planting density. Except at the seeding stage, the biomass accumulation increased rapidly as planting density increased to 12.75 plants/m² (D5), followed by only slight increase with planting density (Fig. 2). However, higher planting density had an adverse effect on the biomass per plant in all seasons (Fig. 3). The biomass accumulation difference and biomass per plant difference between the lowest and highest planting densities reached the maximum at the maturity stage. For example, compared to the lowest planting density (D1), the highest planting density increased biomass accumulation by 55% in 2012, 67% in 2013, 60% in 2014 and 67% in 2015, but decreased biomass per plant by 18% in 2012, 21% in 2013, 24% in 2014 and 21% in 2015, respectively. This can be explained by the fact that the aboveground biomass accumulation

was associated with LAI which increased with planting density. A higher LAI increased the proportion of available radiation interception which then led to higher canopy photosynthesis and biomass accumulation (Loomis and Connor, 1992; Coetto et al., 2013). However, higher planting density resulted in more interplant competition, and thus led to reduced biomass per plant. The above results are consistent with previous findings (Akmal et al., 2014; Thimmappa et al., 2014; Dai et al., 2015).




Fig. 2 Aboveground biomass accumulation of maize for seed production under different planting densities. Error bars denote standard error of the mean.




Fig. 3 Biomass per plant of maize for seed production under different planting densities. Error bars denote standard error of the mean.

3.2 Effect of planting density on yield and WUE

Table 2 presents the yield and WUE results of maize for seed production under different plant densities from 2012 to 2015. It was found that the results showed a large difference in the study years due to the difference in microclimate conditions. The yield per plant and grain yield in 2012 were higher than in other years, while the yield per plant declined with planting density in all the study years. Similar results were also found by Griesh and Yakout (2001) and El-Hendawy et al. (2008). This result might be attributed to more interplant competition at high planting density which led to reduced biomass per plant (Fig. 3) and thus a decreased yield per plant (Kamel et al., 1983; Soliman et al., 1995).

However, the grain yield illustrated different variation trend from yield per plant. It increased rapidly as the planting density increased to 11.25 plants/m² (D4), followed by a slight increase when the planting density was above 11.25 plants/m². The yield started to decrease when the planting density was above 12.75 plants/m². This can be explained by the fact that under lower planting density (D1–D4), the interplant competition was small and thus the grain yield increased with the radiation interception by leaf area which increased during reproductive growth (Johnson et al., 1982; Duncan, 1986). However, when the planting density was great enough to intercept essentially all of the radiation at full canopy, further increase in planting density will not increase the yield (Duncan, 1986). Moreover, a higher planting density may lead to the formation of fewer flower initials, poor pollination resulting from asynchrony of tasseling and silking, or the abortion of kernels after fertilization, and consequently, a reduction of grain yield (Karlen and Camp, 1985; Hashemi-Dezfouli and Herbert, 1992).

The WUE ranged from 1.84 to 2.46 kg/m³ in 2012, 0.90 to 1.19 kg/m³ in 2013, 1.13 to 1.39 kg/m³ in 2014 and 0.86 to 1.36 kg/m³ in 2015, respectively (Table 2). The WUE in 2012 was higher

Table 2 Yield and water use efficiency (WUE) of maize for seed production under different plant densities

Year	Treatment	Yield per plant (g/plant)	Grain yield (t/hm ²)	ET (mm)	WUE (kg/m ³)
2012	D1	121.26 ^a	8.02 ^d	435.74 ^e	1.84 ^d
	D2	116.76 ^{ab}	9.77 ^c	453.43 ^d	2.15 ^c
	D3	111.98 ^{bc}	11.09 ^b	473.97 ^e	2.34 ^b
	D4	106.95 ^{cd}	12.22 ^a	495.68 ^b	2.46 ^a
	D5	100.90 ^d	12.66 ^a	515.67 ^a	2.46 ^a
2013	D1	56.40 ^a	3.81 ^d	421.80 ^f	0.90 ^c
	D2	54.20 ^{ab}	4.47 ^c	443.30 ^e	1.01 ^c
	D3	53.10 ^{ab}	5.17 ^b	462.80 ^d	1.12 ^b
	D4	50.50 ^{bc}	5.68 ^{ab}	486.10 ^c	1.17 ^a
	D5	47.50 ^c	6.05 ^a	507.10 ^b	1.19 ^a
	D6	38.80 ^d	5.53 ^{ab}	531.90 ^a	1.04 ^c
2014	D1	71.23 ^a	4.81 ^d	424.08 ^f	1.13 ^d
	D2	67.32 ^b	5.55 ^c	452.82 ^e	1.23 ^c
	D3	64.84 ^c	6.32 ^b	473.72 ^d	1.33 ^b
	D4	62.40 ^d	7.02 ^a	503.60 ^c	1.39 ^a
	D5	55.55 ^e	7.08 ^a	519.79 ^b	1.36 ^a
	D6	49.30 ^f	7.02 ^a	542.79 ^a	1.29 ^b
2015	D1	73.20 ^a	4.12 ^e	477.00 ^f	0.86 ^d
	D2	71.85 ^a	4.94 ^d	485.35 ^e	1.02 ^c
	D3	69.64 ^b	5.66 ^c	489.53 ^d	1.16 ^b
	D4	68.18 ^b	6.39 ^b	495.29 ^c	1.29 ^a
	D5	63.96 ^c	6.80 ^a	501.12 ^b	1.36 ^a
	D6	47.53 ^d	5.64 ^c	505.83 ^a	1.11 ^b

Note: Different lowercase letters indicate significant differences among different treatments for the same year at the $P < 0.05$ level. ET, evapotranspiration; WUE, water use efficiency.

due to higher grain yield. However, the impact of planting density on WUE was similar for all the study periods. The WUE increased significantly with planting density when it was below 11.25 plants/m² (D4), but no significant increase was observed when the planting density was beyond D4. Table 2 showed that significant decrease of grain yield and WUE was observed when the planting density was below 11.25 plants/m² or above 12.75 plants/m². Thus the densities of 11.25 or 12.75 plants/m² were optimal under the conditions of this study. Cox and Otis (1993) reported the maximum grain yield of maize at 7.41 plants/m². Tetio-Kagho et al. (1998) noted that the grain yield of maize increased parabolically to the maximum yield of 1080 g/m² at about 10.00 plants/m². Sangoi et al. (2002) reported an estimated optimum plant density of maize at about 8.50 plants/m². El-Hendawy et al. (2008) indicated that the optimal density of maize in Egypt was 4.80 or 7.10 plants/m². The optimal planting density in our study was significantly higher than that of hybrid maize. The main reason was that the parent lines of maize for seed production were different from hybrid maize since the female and male parents were short inbred lines without heterosis, and the smaller interplant competition of parent lines led to more tolerance of higher planting density. When the planting density was 11.25 plants/m², WUE reached the maximum. However, the WUE did not significantly increase when the planting density was too high. Moreover, a higher planting density resulted in large evapotranspiration and LAI (Table 2; Fig. 1), leading to the waste of water resources. It also becomes more difficult in efficiently carrying out agronomic practice under a higher planting density. Therefore, it is more economical to select the recommended planting density of 11.25 plants/m² for maize for seed production in Northwest China.

3.3 The modification of yield model under different planting densities

Figure 4 presents the estimated biomass accumulation and grain yield under different planting densities using the AquaCrop yield model (B_A and Y_A) and the modified yield model. Compared to the measured biomass accumulation (B) and grain yield (Y), the results (B_A and Y_A) from AquaCrop yield model were higher than the observed values when the planting density was below the reference planting density, but lower than the observation when the planting density was above the reference planting density (Fig. 4). The determination coefficient (R^2), MAE and RMSE was 0.78, 1.41 and 1.85 t/hm² for biomass accumulation, and 0.88, 0.62 and 0.50 t/hm² for grain yield, respectively (Table 3). The main reason is that the WP* and HI for maize under different planting




Fig. 4 Comparison of modeling results with the measured value B and Y in 2012–2015. Y_A , yield estimated by the AquaCrop yield model; Y_d , yield estimated by the modified AquaCrop yield model; B_A , biomass accumulation estimated by the AquaCrop yield model; B_d , biomass accumulation estimated by the modified AquaCrop yield model.

densities were taken as constants in the AquaCrop yield model, which resulted in large error in model estimation. In fact, they were not constants. For example, WP^* increased with planting density until the optimal density as shown in Figure 5. Moreover, HI increased with planting density when it was below the reference planting density (Fig. 5), but started to decline when the planting density was above the optimum planting density. Similar results were also found in other studies (DeLougherty and Crookston, 1979; Rahmati, 2009). In this study, the effect of planting density on WP^* and HI was considered in the modified AquaCrop yield model. The estimated biomass accumulation and grain yield by the modified model were closer to the observed values (Fig. 4), with higher R^2 , lower MAE and RMSE as compared to the original AquaCrop model (Table 3). As a result, the modified AquaCrop yield model was considered to have a better performance in simulating biomass accumulation and grain yield under different planting densities of maize for seed production.




Fig. 5 Parameter variation of maize for seed production under different planting densities in 2012–2015. WP^* , normalized biomass water productivity; HI, harvest index.

Table 3 Statistical analysis of modeling results from AquaCrop yield model (B_A and Y_A) and modified yield model (B_d and Y_d) in comparison with the measured values (B and Y)

Item	Regression equation	R^2	MAE (t/hm ²)	RMSE (t/hm ²)
Biomass accumulation	$B_A=1.04B$	0.78	1.41	1.85
	$B_d=1.00B$	0.98	0.47	0.56
Grain yield	$Y_A=1.03Y$	0.88	0.62	0.50
	$Y_d=0.97Y$	0.97	0.33	0.21

Note: R^2 , determination coefficient; MAE, mean absolute bias error; RMSE, root mean square error.

4 Conclusions

A field experiment was conducted in 2012–2015 to examine the impacts of planting density on the biomass accumulation and yield of maize for seed production in Northwest China, and the AquaCrop yield model was modified by taking into account the effect of planting density on the WP^* and HI. The results indicated that increasing the planting density led to the increase of LAI and aboveground biomass accumulation, but significantly decreased the yield and biomass per plant. The grain yield and WUE increased rapidly as planting density increased to 11.25 plants/m², but there was only a slight increase in the grain yield when the planting density was above 11.25 plants/m². It is thus more economical to select a recommended planting density of 11.25 plants/m² for maize for seed production in Northwest China. Within the range of planting densities used in the experiments, the original AquaCrop yield model produced a large error, with significantly greater estimation than the measured biomass accumulation and grain yield when the planting density was below the reference planting density. It had a lower estimation than the measured value when the planting density was above the reference planting density. However, the estimated results using the modified yield model were closer to the measured values, with higher determination

coefficient, and lower MAE and RMSE, indicating that the modified yield model by considering the effect of planting density had a better performance in estimating the biomass accumulation and grain yield of maize for seed production in the arid region of Northwest China.

Acknowledgements

We are grateful for the research grants from the National Natural Science Foundation of China (51379208, 91425302, 51621061), the Government Public Research Funds for Projects of the Ministry of Agriculture (201503125) and the Discipline Innovative Engineering Plan (111 Program, B14002).

References

- Akmal M, Asim M, Gilbert M. 2014. Influence of seasonal variation on radiation use efficiency and crop growth of maize planted at various densities and nitrogen rates. *Pakistan Journal of Agricultural Sciences*, 51(4): 835–846.
- Allen R G, Pereira L S, Raes D, et al. 1998. Crop evaporation: guidelines for computing crop water requirements. In: *FAO Irrigation and Drainage Paper No. 56*. Rome: FAO, 300.
- Coetto E, Di Candillo M, Castelli F, et al. 2013. Comparing solar radiation interception and use efficiency for the energy crops giant reed (*Arundo donax* L.) and sweet sorghum (*Sorghum bicolor* L. Moench). *Field Crops Research*, 149: 159–166.
- Cox W J, Otis D J. 1993. Grain and silage yield responses of commercial corn hybrids to plant densities. In: *Agronomy Abstracts*. Madison, Wisconsin: American Society of Agronomy, 132.
- Dai J L, Li W J, Tang W, et al. 2015. Manipulation of dry matter accumulation and partitioning with plant density in relation to yield stability of cotton under intensive management. *Field Crops Research*, 180: 207–215.
- DeLougherty R L, Crookston R K. 1979. Harvest index of corn affected by population density, maturity rating, and environment. *Agronomy Journal*, 71(4): 577–580.
- Duncan W G. 1986. Planting patterns and soybean yields. *Crop Science*, 26(3): 584–588.
- El-Hendawy S E, El-Lattief E A A, Ahmed M S, et al. 2008. Irrigation rate and plant density effects on yield and water use efficiency of drip-irrigated corn. *Agricultural Water Management*, 95(7): 836–844.
- Fulton J M. 1970. Relationships among soil moisture stress, plant populations, row spacing and yield of corn. *Canadian Journal of Plant Science*, 50(1): 31–38.
- Griesh M H, Yakout G M. 2001. Effect of plant population density and nitrogen fertilization on yield and yield components of some white and yellow maize hybrids under drip irrigation system in sandy soil. In: Horst W J, Schenk M K, Bürkert A, et al. *Plant Nutrition*. Dordrecht: Springer, 810–811.
- Harper L A, Pallas J E Jr, Bruce R R, et al. 1979. Greenhouse microclimate for tomatoes in the southeast USA. *American Society for Horticultural Science*, 104(5): 659–663.
- Hashemi-Dezfouli A, Herbert S J. 1992. Intensifying plant density response of corn with artificial shade. *Agronomy Journal*, 84(4): 547–551.
- Holt D F, Timmons D R. 1968. Influence of precipitation, soil water, and plant population interactions on corn grain yields. *Agronomy Journal*, 60(4): 379–381.
- Jiang X L, Kang S Z, Tong L, et al. 2014. Crop coefficient and evapotranspiration of grain maize modified by planting density in an arid region of northwest China. *Agricultural Water Management*, 142: 135–143.
- Johnson R R, Green D E, Jordan C W. 1982. What is the best soybean row width? A U.S. perspective. *Crops and Soils Magazine*, 43(4): 10–13.
- Kamel M S, El-Raouf M S, Mahmoud E A, et al. 1983. Response of two maize varieties to different plant densities in relation to weed control treatments. *Annals of Agricultural Sciences*, 19: 79–93.
- Karlen D L, Camp C R. 1985. Row spacing, plant population, and water management effects on corn in the Atlantic Coastal Plain. *Agronomy Journal*, 77(3): 393–398.
- Lang A L, Pendleton J W, Dungan G H. 1956. Influence of population and nitrogen levels on yield and protein and oil contents of nine corn hybrids. *Agronomy Journal*, 48(7): 284–289.
- Legates D R, McCabe G J Jr. 1999. Evaluating the use of “goodness-of-fit” measures in hydrologic and hydroclimatic model validation. *Water Resources Research*, 35(1): 233–241.
- Lemcoff J H, Loomis R S. 1986. Nitrogen influences on yield determination in maize. *Crop Science*, 26(5): 1017–1022.
- Li S E, Kang S Z, Li F S, et al. 2008. Evapotranspiration and crop coefficient of spring maize with plastic mulch using eddy covariance in northwest China. *Agricultural Water Management*, 95(11): 1214–1222.

- Loomis R S, Connor D J. 1992. *Crop Ecology: Productivity and Management in Agricultural Systems*. Cambridge, UK: Cambridge University Press.
- Mohamed M M A. 1999. Effect of some agronomic practices on corn production (*Zea mays* L.) under drip irrigation system. PhD Dissertation. Ismailia, Egypt: Faculty of Agriculture, Suez Canal University, 107.
- Olson R A, Sander D H. 1988. Corn production. In: Sprague G F, Dudley J W. *Corn and Corn Improvement* (3rd ed.). Madison, WI: American Society of Agronomy, 639–686.
- Papadopoulos A P, Pararajasingham S. 1997. The influence of plant spacing on light interception and use in greenhouse tomato (*Lycopersicon esculentum* Mill.): a review. *Scientia Horticulturae*, 69(1–2): 1–29.
- Qiu R J, Song J J, Du T S, et al. 2013. Response of evapotranspiration and yield to planting density of solar greenhouse grown tomato in northwest China. *Agricultural Water Management*, 130: 44–51.
- Raes D, Steduto P, Hsiao T C, et al. 2009. AquaCrop—The FAO crop model to simulate yield response to water: II. Main algorithms and software description. *Agronomy Journal*, 101(3): 438–477.
- Raes D, Steduto P, Hsiao C, et al. 2011. *AquaCrop Version 3.1Plus Reference Manual*. Rome, Italy: FAO.
- Rahmati H. 2009. Effect of plant density and nitrogen rates on yield and nitrogen efficiency of grain corn. *World Applied Science Journal*, 7(8): 958–961.
- Rana G, Katerji N. 2000. Measurement and estimation of actual evapotranspiration in the field under Mediterranean climate: a review. *European Journal of Agronomy*, 13(2–3): 125–153.
- Sangoi L, Gracietti M A, Rampazzo C, et al. 2002. Response of Brazilian maize hybrids from different eras to changes in plant density. *Field Crops Research*, 79(1): 39–51.
- Soliman F H, Goda A S, Ragheb M M, et al. 1995. Response of maize (*Zea mays* L.) hybrids to plant populations density under different environmental conditions. *Hip International*, 23(6):124–124.
- Steduto P, Hsiao T C, Raes D, et al. 2009. AquaCrop—the FAO crop model to simulate yield response to water: I. Concepts and underlying principles. *Agronomy Journal*, 101(3): 426–437.
- Tetio-Kagho F, Gardner F P. 1998. Responses of maize to plant population density. II. Reproductive development, yield, and yield adjustments. *Agronomy Journal*, 80(6): 935–940.
- Thimmappa V, Reddy M S, Reddy U, et al. 2014. Effect of nitrogen levels and plant densities on growth parameters, yield attributes and yield of kharif maize (*Zea mays* L.). *Crop Research*, 47(1–3): 29–32.
- Westgate M E, Forcella F, Reicosky D C, et al. 1997. Rapid canopy closure for maize production in the northern US corn belt: Radiation-use efficiency and grain yield. *Field Crops Research*, 49(2–3): 249–258.